

Betacol®

Betacol Promotes Healthy Liver, Cardiovascular, and Cellular Function

Betacol contains *Tillandsia usneoides*, a source of important phytochemicals that contribute to cardiovascular and cellular health. Several bioactive compounds, including vitamin C, carotene, and HMG (3-hydroxy-3-methylglutaric acid), are found in *Tillandsia usneoides*. Vitamins, minerals, and phytochemicals in Betacol work at the cellular level to maintain healthy blood glucose and cholesterol levels in individuals with normal levels. Betaine in Betacol is important for healthy liver function, especially the metabolism of fat.†

How Betacol Keeps You Healthy

Promotes healthy liver function

Betaine is a lipotropic agent that promotes transportation and metabolism of fats, helping to prevent the accumulation of fat in the liver. Several studies demonstrate betaine's unique ability to support a healthy liver.†

Supports cardiovascular health

Tillandsia usneoides supports arterial health by helping to maintain normal serum cholesterol levels in individuals with normal levels. HMG, the bioactive compound in *Tillandsia usneoides*, has been shown in several studies to help support cardiovascular health factors. Several studies have confirmed that HMG operates at the enzymatic level in the kidney and intestine. Studies also indicate that HMG may help maintain healthy cardiovascular function. Betacol contains vitamin B₆, which is important for red blood cell formation and helps maintain optimum homocysteine levels in the blood. Vitamin B₆ is well documented for its ability to support the heart muscle and associated blood vessels.†

Promotes healthy cellular function

Vitamin B₆ is required to synthesize the nucleic acids RNA and DNA—the molecules that carry the genetic instructions for normal cellular growth and reproduction. Betacol also contains niacin, which functions as a coenzyme that is key to cellular respiration, carbohydrate and protein metabolism, and lipid synthesis. Niacin is a hydrogen acceptor, combining with hydrogen atoms as they are removed from the food we eat. Once this occurs, the coenzyme continues transferring these atoms to other compounds in a series of complex oxidation reactions. Several studies indicate that extracts from *Tillandsia usneoides* contain compounds capable of reducing serum glucose. HMG may also help support liver function.†



Introduced in: 1946

Content: 40 Capsules

Supplement Facts:

Serving Size: 1 capsule		
Servings per Container: 40		
		%DV
Calories	2	
Niacin	10 mg	50%
Vitamin B ₆	2 mg	100%

Proprietary Blend: *Tillandsia usneoides*, soy (bean), bovine liver PMG™ extract, betaine hydrochloride, calcium lactate, defatted wheat (germ), potassium bicarbonate, choline bitartrate, inositol, disodium phosphate, bovine adrenal Cytosol™ extract, oat flour, and ascorbic acid.

Other Ingredients: Gelatin, water, niacinamide, pyridoxine hydrochloride, colors, and calcium stearate.

Suggested Use: One capsule per meal, or as directed.

Sold through health care professionals.

Please call for your patients.



800-558-8740 | www.standardprocess.com

Betacol®

Betacol[®]

What Makes Betacol Unique

Product Attributes

A distinctive product that contains choline

- ▶ Choline is essential to metabolize fat cholesterol, proteins, and carbohydrates effectively†

Multiple nutrients from a variety of plant and animal sources

- ▶ Extracts from bovine tissues provide nutrients and support to the corresponding tissues in humans
- ▶ Vitamins, minerals, and nutrients from plants and animal tissues work synergistically for maximum effect†

Contains Protomorphogen™ extracts

- ▶ Standard Process uses a unique manufacturing method of deriving tissue cell determinants from animal glands and organs
- ▶ Help provide cellular support and rehabilitation to the corresponding human tissues
- ▶ Important antigenic properties of nucleoprotein-mineral determinants are the foundation of the product†

Manufacturing and Quality Control Processes

Low-temperature, high-vacuum drying technique

- ▶ Preserves the enzymatic vitality and nutritional potential of ingredients

Degreed microbiologists and chemists in our on-site laboratories continually conduct bacterial and analytical tests on raw materials, product batches, and finished products

- ▶ Ensures consistent quality and safety

Vitamin and mineral analyses validate product content and specifications

- ▶ Assures high-quality essential nutrients are delivered

Whole Food Philosophy

Our founder, Dr. Royal Lee challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists—known and unknown—bioactivity is markedly enhanced over isolated nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to an isolated or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

Statistics on nutrients generally use large doses and these studies, some of which are cited below, are the cases for much of the information we provide you in the publication about whole food ingredients. See the supplement facts for Betacol.

- Anderson L.E. 1999. *Wiley's Medical, Nursing & Allied Health Dictionary*, 6th ed. St. Louis, MO: Wiley; 1999.
- Ang M.P. *Spanish Moss and Sphagnum*. University of Florida, Cooperative Extension Service Institute of Food and Agricultural Sciences. Online 21 May 2000.
- Baker J.R., Bach-Peterson A. 1992. *Prescription for Additional Healing*, 2nd ed. Garden City Park, NY: Avery Publishing Group; 19-19.
- Bardner C.D. 1995. *Advanced Nutrition*. Westborough: Boca Raton, FL: CRC Press; 64-109.
- Coffin C.J. 1998. *Metabolic Inhibitors*. Madison, WI: Perseus Books Publishing; 69.
- Costa M., DiStasio L.C., Krasner M., et al. 1998. Screening in mice of some medicinal plants used for weight loss purposes in the state of São Paulo, part II. *J Ethnopharmacol* 27(1-2): 25-33.
- Di Paolo G., Bazzani E., Ughetti R., et al. 1992. 3-hydroxy-3-methylglutaryl coenzyme A (HMG) reductase activity and inhibition of statins in human liver. *J Lipid Res* 33(22): 3517-3524.
- DuBo J. USDA-ARS-NRRL. Phytochemistry and Ethnobotanical Database. Boca Raton, FL: Boca Raton Agricultural Research Center. Online 22 May 2000.
- Guyton A.C., Hall J.E. 1997. *Human Physiology and Mechanism of Disease*, 6th ed. Philadelphia, PA: WB Saunders Company; 692.
- Kraschinsky J.D. 1979. *Natural Alkaloids*. Revised ed. New York, NY: McGraw-Hill Book Company; 25-27.
- Labate J.E., Dam R. 1960. Effect of 3-hydroxy-3-methylglutaryl coenzyme A reductase in female Japanese quail. *Proc Soc Exp Biol Med* 105: 385-386.
- Lupien P.J., Morgan S., Bui D., et al. 1970. Effects of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors on cholesterol levels in female hypercholesterolemia. *J Clin Pharmacol* 10(2-3): 120-125.
- Morgan S., Lupien P.J. 1977. Effect of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors on the synthesis of mevalonate acid and its precursors. *Arch Int Physiol Biochem* 85(1): 1-10.
- Murakami T., Nagamura Y., Hara K. 1969. The reducing effect of basane on carbon tetrachloride-induced liver injury. *J Nutr Sci Vitaminol* 44(2): 249-255.
- Pritchard P. 1969. *Healing With Whole Foods*. Revised ed. Berkeley, CA: North Atlantic Books; 122, 402-403.
- Saizumi L., Lupien P.J. 1975. Cogen distribution of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor. *Can J Physiol Pharmacol* 53(4): 638-643.
- Saizumi L., Lupien P.J. 1976. Preliminary toxicological investigations of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor. *J Natl Sci* 1284-1285.
- Shils M.E., Young V.R. 1985. *Nutrient Metabolism in Health and Disease*, 7th ed. Philadelphia, PA: Lea and Febiger; 370-381.
- Winters J. *Winters' Ecology and Evolutionary Biology*. Conservancy Online 17 May 2000.
- Ward D.F., Russell P. 1959. *The Nutrition and Health Encyclopedia*, 2nd ed. New York, NY: Van Nostrand Reinhold; 358-359, 445-449.
- Van Nieuwenhuijzen, J., Harshbarger T., Sauer-Lewis J.A., et al. 1984. The pharmacological effects of 3-hydroxy-3-methylglutaryl coenzyme A and choline in WHHL rabbits. *Clin Sci* 67(4): 439-444.
- Wilson E.O., Fester K.H., Fujita M.E. 1965. *Prosopis juliflora* and related species. *NY State Agric & Mech Expt Sta* 250-254.
- Wolfe A.P., McLaughlin J.L., Judd R.L., et al. 1995. Identification of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor principle of Spanish moss (*Wendlandia bartramia*). *J Natl Prod* 58(8): 1285-1290.
- Yoshida S.Y., Siddle M. 1976. 3-Hydroxy-3-methylglutaryl coenzyme A reductase inhibitors in rats. *Biochem J* 156: 1179-1179.
- Yoshida S.Y., Siddle M. 1977. Saturation and lipid composition of 3-hydroxy-3-methylglutaryl coenzyme A reductase in different carbohydrate diets. *Lipids* 12(3): 262-265.

†These statements have not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

